



Patient's name:

Doctor's name:

Doctor's phone:

For further information, please talk to your health professional, visit **medsafe.govt.nz** for OCREVUS Consumer Medicine Information or download the OCREVUS Patient Booklet by scanning the QR code or visiting **getonwithlife.co.nz**



Roche Products (New Zealand) Limited Auckland
Medical Information Ph 0800 276 243
www.roche.co.nz

M-NZ-00001424/MR12421/SEP2025 ROC00849

OCREVUS[®]
ocrelizumab
Subcutaneous injection



Patient Information Card

This card contains important safety information that you need to be aware of during treatment with OCREVUS subcutaneous (SC).

Injection reactions may occur during the OCREVUS SC injection and **up to 24 hours later.**

Tell your doctor immediately or go to Accident and Emergency at your nearest hospital if you notice any of the following:

- swelling of your face, lips, tongue or throat with difficulty breathing
- swelling of other parts of your body
- shortness of breath, wheezing or trouble breathing
- rash, itching or hives on the skin
- feeling sick (nausea)
- cough, throat irritation or pain
- fever, flushing or chills
- feeling tired
- headache
- dizziness or light headedness
- chest pain
- abnormal or irregular heartbeat



Report any signs of being unwell.

Infections: OCREVUS SC increases your risk of getting an infection. Notify your doctor or nurse right away if you develop signs of an infection such as fever and/or chills, constant cough or signs of herpes (cold sores, shingles or genital sores).

Changes in MS: Tell your partner or caregiver you are receiving OCREVUS SC and ask them to tell you if they notice any changes in your movement or behaviour. If they notice any changes you should tell your doctor or nurse about them immediately.