

Roche

Patient's name:

Doctor's name:

Doctor's phone:

For further information, please talk to your health professional, visit **medsafe.govt.nz** for OCREVUS Consumer Medicine Information or download the OCREVUS Patient Booklet by scanning the QR code or visiting **getonwithlife.co.nz**



Roche Products (New Zealand) Limited Auckland
Medical Information Ph 0800 276 243
www.roche.co.nz

M-NZ-00001424v3/DA2521KS/JUL2026

OCREVUS[®]
ocrelizumab
Subcutaneous injection



Patient Information Card

This card contains important safety information that you need to be aware of during treatment with OCREVUS subcutaneous (SC).

Injection reactions may occur during the OCREVUS SC injection and **up to 24 hours later.**

Tell your doctor immediately or go to Accident and Emergency at your nearest hospital if you notice any of the following:

- itchy skin, rash, hives or redness of the skin
- throat irritation or pain
- shortness of breath
- swelling of the throat, face, lips, tongue, or other parts of the body
- reactions in the area where the injection is given (OCREVUS SC only)
- redness on your face (flushing)
- low blood pressure
- fever
- fatigue
- nausea
- headache
- fast heart beat



Report any signs of being unwell.

Infections: OCREVUS SC increases your risk of getting an infection. Notify your doctor or nurse right away if you develop signs of an infection such as fever and/or chills, constant cough or signs of herpes (cold sores, shingles or genital sores), or COVID-19 pneumonia.

Changes in MS: Tell your partner or caregiver you are receiving OCREVUS SC and ask them to tell you if they notice any changes in your movement or behaviour. If they notice any changes you should tell your doctor or nurse about them immediately.

Liver problems: Tell your doctor if you have any signs and symptoms that may indicate a liver problem (e.g. new or worsening fatigue, loss of appetite, nausea, vomiting, pain in the right part of your abdomen, dark-colored urine, or yellowing of your skin or eyes).