



Patient's name:

Doctor's name:

Doctor's phone:

For further information, please talk to your health professional, visit www.medsafe.govt.nz for OCREVUS Consumer Medicine Information or download the OCREVUS Patient Booklet at www.getonwithlife.co.nz

Roche Products (New Zealand) Limited Auckland
Medical Information Ph 0800 276 243
www.roche.co.nz

M-NZ-00000066-V2.0/MR9351/Jun2023 ROC00711



Anti-CD20 medicine for multiple sclerosis

Patient Information Card

This card contains important safety information that you need to be aware of during treatment with OCREVUS.



Infusion-related reactions may occur during the infusion of OCREVUS and **up to 24 hours later.**

Tell your doctor immediately or go to Accident and Emergency at your nearest hospital if you notice any of the following:

- swelling of your face, lips, tongue or throat with difficulty breathing
- swelling of other parts of your body
- shortness of breath, wheezing or trouble breathing
- skin problems including rash, itchiness or hives
- fever, flushing or chills
- cough, throat irritation or pain
- feeling tired
- headache
- dizziness or light headedness



Report any signs of being unwell.

- **Infections:** OCREVUS increases your risk of getting an infection. Notify your doctor or nurse right away if you develop signs of an infection such as fever and/or chills, constant cough or signs of herpes (cold sores, shingles or genital sores).
- **Changes in MS:** Tell your partner or caregiver you are receiving OCREVUS and ask them to tell you if they notice any changes in your movement or behaviour. If they notice any changes you should tell your doctor or nurse about them immediately.